


HIV Surveillance by Race/Ethnicity

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention



Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity, 2008–2011


United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Rates of Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity 2008–2011—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.

HIV Infection in Blacks/African Americans United States and 6 Dependent Areas

Of the 197,090 diagnoses of HIV infection from 2008–2011,
blacks/African Americans accounted for:

- 47% of total

- 64% of women

- 66% of infections attributed to heterosexual contact^a

- 67% of children aged <13 years

In 2011, 46% of diagnoses of HIV infection among adults and
adolescents were in blacks/African Americans.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



HIV Infection in Hispanics/Latinos United States and 6 Dependent Areas

Of the 197,090 diagnoses of HIV infection from 2008–2011, Hispanics/Latinos^a accounted for:

- 21% of total

- 16% of women

- 18% of infections attributed to heterosexual contact^b

- 15% of children aged <13 years

In 2011, 22% of diagnoses of HIV infection among adults and adolescents were in Hispanics/Latinos.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting

^a Hispanics/Latinos can be of any race.


^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



Diagnoses of HIV Infection and Population by Race/Ethnicity, 2011—United States

Diagnoses of HIV infection


N = 49,273



■ American Indian/Alaska Native
■ Asian
■ Black/African American

Population, United States

N = 311,591,917



■ Hispanic/Latino^a
■ Native Hawaiian/other Pacific Islander
■ White


■ Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.


Diagnoses of HIV Infection and Population among Adult and Adolescent Males, by Race/Ethnicity 2011—United States

Diagnoses of HIV Infection
N = 38,825



■ American Indian/Alaska Native
■ Asian
■ Black/African American

Male Population, United States
N = 126,238,952



■ Hispanic/Latino^a
■ Native Hawaiian/Other Pacific Islander
■ White

■ Multiple races


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection and Population among Adult and Adolescent Females, by Race/Ethnicity 2011—United States

Diagnoses of HIV Infection

N = 10,257




American Indian/Alaska Native
Asian
Black/African American

Multiple races

Female Population, United States

N = 132,402,857




Hispanic/Latino^a
Native Hawaiian/other Pacific Islander
White


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, by Sex and Transmission Category 2011—United States and 6 Dependent Areas



- Male-to-male sexual contact
- Injection drug use
- Male-to-male sexual contact and IDU





- Heterosexual contact^a
- Other^b

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, by Sex and Transmission Category 2011—United States and 6 Dependent Areas



- Male-to-male sexual contact
- Injection drug use
- Male-to-male sexual contact and IDU

- Heterosexual contact^b
- Other^c


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.


^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Diagnoses of HIV Infection among Adult and Adolescent Whites, by Sex and Transmission Category 2011—United States and 6 Dependent Areas



- Male-to-male sexual contact
- Injection drug use
- Male-to-male sexual contact and IDU



- Heterosexual contact^a
- Other^b

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, by Sex and Place of Birth 2011—United States and 6 Dependent Areas


Males

N = 9,256



Females

N = 1,776



- United States
- Central America
- South America
- Cuba
- Mexico
- Puerto Rico
- Other
- Unknown

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection among Adult and Adolescent Males, by Race/Ethnicity, 2011—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	161	18.0
Asian	821	13.8
Black/African American	16,447	112.8
Hispanic/Latino ^a	8,605	43.4
Native Hawaiian/other Pacific Islander	70	34.2
White	12,041	14.5
Multiple races	679	38.5
Total	38,825	30.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2011—United States


Race/ethnicity	No.	Rate
American Indian/Alaska Native	51	5.5
Asian	153	2.3
Black/African American	6,595	40.0
Hispanic/Latino ^a	1,530	7.9
Native Hawaiian/Other Pacific Islander	8	3.9
White	1,776	2.0
Multiple races	144	7.5
Total	10,257	7.7

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adults and Adolescents, by Region and Race/Ethnicity 2011—United States



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.


^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, 2011—United States

N = 23,042

Total Rate = 74.2




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Rates of Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, 2011—United States

N = 10,134

Total Rate = 25.9



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Whites, 2011—United States

N = 13,817

Total Rate = 8.1



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Deaths of Persons with Diagnosed HIV Infection by Race/Ethnicity, 2010—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	90	3.9
Asian ^a	91	0.6
Black/African American	9,517	25.0
Hispanic/Latino ^b	2,846	5.6
Native Hawaiian/other Pacific Islander	10	2.1
White	5,981	3.0
Multiple races	803	14.2
Total^c	19,343	6.3

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.



Adults and Adolescents Living with Diagnosed HIV Infection by Race/Ethnicity, Year-end 2010—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	3,187	176.4
Asian ^a	9,646	77.8
Black/African American	381,399	1,242.4
Hispanic/Latino ^b	164,654	432.4
Native Hawaiian/other Pacific Islander	892	223.6
White	295,146	174.2
Multiple races	14,412	404.7
Total^c	870,096	339.4

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.




Stage 3 (AIDS) Classifications among Racial/Ethnic Minorities with Diagnosed HIV Infection, 1985–2011 United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Includes American Indians/Alaska Natives, Asians, blacks/African Americans, Hispanics/Latinos, and Native Hawaiians/other Pacific Islanders. Hispanics/Latinos can be of any race.

Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity, 1985–2011 United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.

Rates of Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity 2000–2011—United States



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.


^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications and Population, by Race/Ethnicity, 2011—United States

Stage 3 (AIDS)

N = 32,052




■ American Indian/Alaska Native
■ Asian^b
■ Black/African American

■ Multiple races

U.S. Population

N = 311,591,917



■ Hispanic/Latino^a
■ Native Hawaiian/other Pacific Islander
■ White

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.

Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Blacks/African Americans with Diagnosed HIV Infection, 2011—United States

N = 15,946

Total Rate = 51.3




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.

Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Hispanics/Latinos^a with Diagnosed HIV Infection, 2011—United States

N = 6,353

Total Rate = 16.2



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.

^aHispanics/Latinos can be of any race.




Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Whites with Diagnosed HIV Infection, 2011

United States


N = 8,303

Total Rate = 4.9



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.

Survival after Classification of Stage 3 (AIDS) during 1998–2007, by Months Survived and Race/Ethnicity United States and 6 Dependent Areas



Note. Data exclude persons whose month of diagnosis or month of death is unknown.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, Year-end 2010—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	1,694	93.7
Asian ^a	5,360	43.2
Black/African American	209,198	681.4
Hispanic/Latino ^b	99,515	261.3
Native Hawaiian/Other Pacific Islander	488	122.2
White	162,208	95.7
Multiple races	8,679	243.7
Total^c	487,235	190.0

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.

